arXiv:1408.1269v1 [math.FA] 6 Aug 2014

# On some new difference sequence spaces of fractional order

Serkan Demiriz*,[a], Osman Duyar[b]

[a]*Department of Mathematics, Faculty of Arts and Science, Gaziosmanpaşa University, 60250 Tokat, Turkey*
[b]*Anatolian High School, 60200 Tokat, Turkey*

**Abstract**

Let $\Delta^{(\alpha)}$ denote the fractional difference operator. In this paper, we define new difference sequence spaces $c_0(\Gamma, \Delta^{(\alpha)}, u)$ and $c(\Gamma, \Delta^{(\alpha)}, u)$. Also, the $\beta-$dual of the spaces $c_0(\Gamma, \Delta^{(\alpha)}, u)$ and $c(\Gamma, \Delta^{(\alpha)}, u)$ are determined and calculated their Schauder basis. Furthermore, the classes $(\mu(\Gamma, \Delta^{(\alpha)}, u) : \lambda)$ where $\mu \in \{c_0, c\}$ and $\lambda \in \{c_0, c, \ell_\infty, \ell_1\}$ .

*Key words:* Difference operator $\Delta^{(\alpha)}$, Sequence spaces, $\beta-$dual, Matrix transformations.

## 1. Preliminaries,background and notation

By a *sequence space*, we mean any vector subspace of $\omega$, the space of all real or complex valued sequences $x = (x_k)$. The well-known sequence spaces that we shall use throughout this paper are as following:

$\ell_\infty$: the space of all bounded sequences,

$c$: the space of all convergent sequences,

$c_0$: the space of all null sequences,

$cs$: the space of all sequences which form convergent series,

$\ell_1$: the space of all sequences which form absolutely convergent series,

$\ell_p$: the space of all sequences which form $p$-absolutely convergent series, where $1 < p < \infty$.

*Corresponding Author (Tel: +90 356 252 16 16, Fax: +90 356 252 15 85)

*Email addresses:* serkan.demiriz@gop.edu.tr (Serkan Demiriz), osman-duyar@hotmail.com (Osman Duyar)

Let $X, Y$ be two sequence spaces and $A = (a_{nk})$ be an infinite matrix of real or complex numbers $a_{nk}$, where $n, k \in \mathbb{N}$. Then, we say that $A$ defines a matrix mapping from $X$ into $Y$, and we denote it by writing $A : X \to Y$, if for every sequence $x = (x_k) \in \lambda$ the sequence $Ax = \{(Ax)_n\}$, the $A$-transform of $x$, is in $Y$; where

$$(Ax)_n = \sum_k a_{nk} x_k, \quad (n \in \mathbb{N}). \tag{1}$$

For simplicity in notation, here and in what follows, the summation without limits runs from 0 to $\infty$. By $(X : Y)$, we denote the class of all matrices $A$ such that $A : X \to Y$. Thus, $A \in (X : Y)$ if and only if the series on the right side of (1) converges for each $n \in \mathbb{N}$ and every $x \in X$, and we have $Ax = \{(Ax)_n\}_{n\in\mathbb{N}} \in Y$ for all $x \in X$. A sequence $x$ is said to be $A$-summable to $\alpha$ if $Ax$ converges to $\alpha$ which is called as the $A$-limit of $x$.

If a normed sequence space $X$ contains a sequence $(b_n)$ with the property that for every $x \in X$ there is a unique sequence of scalars $(\alpha_n)$ such that

$$\lim_{n\to\infty} \|x - (\alpha_0 b_0 + \alpha_1 b_1 + ... + \alpha_n b_n)\| = 0,$$

then $(b_n)$ is called a *Schauder basis* (or briefly *basis*) for $X$. The series $\sum \alpha_k b_k$ which has the sum $x$ is then called the expansion of $x$ with respect to $(b_n)$, and written as $x = \sum \alpha_k b_k$.

A matrix $A = (a_{nk})$ is called a *triangle* if $a_{nk} = 0$ for $k > n$ and $a_{nn} \neq 0$ for all $n \in \mathbb{N}$. It is trivial that $A(Bx) = (AB)x$ holds for triangle matrices $A, B$ and a sequence $x$. Further, a triangle matrix $U$ uniquely has an inverse $U^{-1} = V$ that is also a triangle matrix. Then, $x = U(Vx) = V(Ux)$ holds for all $x \in \omega$. We write additionally $\mathcal{U}$ for the set of all sequences $u$ such that $u_k \neq 0$ for all $k \in \mathbb{N}$.

For a sequence space $X$, the *matrix domain* $X_A$ of an infinite matrix $A$ is defined by

$$X_A = \{x = (x_k) \in \omega : Ax \in \lambda\}, \tag{2}$$

which is a sequence space. The approach constructing a new sequence space by means of the matrix domain of a particular limitation method has been recently employed by Wang [6], Ng and Lee [7], Aydın and Başar [17] and Altay and Başar [12].

The gamma function may be regarded as a generalization of $n!$ ($n-$factorial), where $n$ is any positive integer. The gamma function $\Gamma$ is defined for all $p$

real numbers except the negative integers and zero. It can be expressed as an improper integral as follows:

$$\Gamma(p) = \int_0^\infty e^{-t}t^{p-1}dt. \tag{3}$$

From the equality (3) we deduce following properties:

(i) If $n \in \mathbb{N}$ then we have $\Gamma(n+1) = n!$
(ii) If $n \in \mathbb{R} - \{0, -1, -2, -3...\}$ then we have $\Gamma(n+1) = n\Gamma(n)$.

For a proper fraction $\alpha$, Baliarsingh and Dutta have defined a fractional difference operators $\Delta^\alpha : w \to w$, $\Delta^{(\alpha)} : w \to w$ and their inverse in [20] as follows:

$$\Delta^\alpha(x_k) = \sum_{i=0}^{\infty}(-1)^i\frac{\Gamma(\alpha+1)}{i!\Gamma(\alpha+1-i)}x_{k+i} \tag{4}$$

$$\Delta^{(\alpha)}(x_k) = \sum_{i=0}^{\infty}(-1)^i\frac{\Gamma(\alpha+1)}{i!\Gamma(\alpha+1-i)}x_{k-i} \tag{5}$$

$$\Delta^{-\alpha}(x_k) = \sum_{i=0}^{\infty}(-1)^i\frac{\Gamma(1-\alpha)}{i!\Gamma(1-\alpha-i)}x_{k+i} \tag{6}$$

and

$$\Delta^{(-\alpha)}(x_k) = \sum_{i=0}^{\infty}(-1)^i\frac{\Gamma(1-\alpha)}{i!\Gamma(1-\alpha-i)}x_{k-i} \tag{7}$$

where we assume throughout the series defined in (4)-(7) are convergent. In particular, for $\alpha = \frac{1}{2}$,

$$\begin{aligned}
\Delta^{1/2}x_k &= x_k - \frac{1}{2}x_{k+1} - \frac{1}{8}x_{k+2} - \frac{1}{16}x_{k+3} - \frac{5}{128}x_{k+4} - \frac{7}{256}x_{k+5} - ... \\
\Delta^{-1/2}x_k &= x_k + \frac{1}{2}x_{k+1} + \frac{3}{8}x_{k+2} + \frac{5}{16}x_{k+3} + \frac{35}{128}x_{k+4} + \frac{63}{256}x_{k+5} + ... \\
\Delta^{(1/2)}x_k &= x_k - \frac{1}{2}x_{k-1} - \frac{1}{8}x_{k-2} - \frac{1}{16}x_{k-3} - \frac{5}{128}x_{k-4} - \frac{7}{256}x_{k-5} - ... \\
\Delta^{(-1/2)}x_k &= x_k + \frac{1}{2}x_{k-1} + \frac{3}{8}x_{k-2} + \frac{5}{16}x_{k-3} + \frac{35}{128}x_{k-4} + \frac{63}{256}x_{k-5} + ....
\end{aligned}$$

Baliarsingh have been defined the spaces $X(\Gamma, \Delta^{\alpha}, u)$ for $X \in \{\ell_{\infty}, c_0, c\}$ by introducing the fractional difference operator $\Delta^{\alpha}$ and a positive fraction $\alpha$ in [18]. In this article, Baliarsingh have been studied some topological properties of the spaces $X(\Gamma, \Delta^{\alpha}, u)$ and established their $\alpha-$, $\beta-$ and $\gamma-$duals.

Following [18], we introduce the sequence spaces $c_0(\Gamma, \Delta^{(\alpha)}, u)$ and $c(\Gamma, \Delta^{(\alpha)}, u)$ and obtain some results related to these sequence spaces. Furthermore, we compute the $\beta-$dual of the spaces $c_0(\Gamma, \Delta^{(\alpha)}, u)$ and $c(\Gamma, \Delta^{(\alpha)}, u)$ . Finally, we characterize some matrix transformations on new sequence spaces.

## 2. The sequence spaces $c_0(\Gamma, \Delta^{(\alpha)}, u)$ and $c(\Gamma, \Delta^{(\alpha)}, u)$

In this section, we define the sequence spaces $c_0(\Gamma, \Delta^{(\alpha)}, u)$ and $c(\Gamma, \Delta^{(\alpha)}, u)$ and examine the some topological properties of this sequence spaces.

The notion of difference sequence spaces was introduced by Kızmaz [2]. It was generalized by Et and Çolak [4] as follows:

Let $m$ be a non-negative integer. Then

$$\Delta^m(X) = \{x = (x_k) : \Delta^m x_k \in X\}$$

where $\Delta^0 x = (x_k)$, $\Delta^m x = (\Delta^{m-1} x_k - \Delta^{m-1} x_{k+1})$ for all $k \in \mathbb{N}$ and

$$\Delta^m x_k = \sum_{i=0}^{m} (-1)^i \binom{m}{i} x_{k+i}. \tag{8}$$

Furthermore, Malkowsky E., et al.[9] have been introduced the spaces

$$\Delta_u^{(m)} X = \{x \in \omega : \Delta_u^{(m)} x \in X\} \tag{9}$$

where $\Delta_u^{(m)} x = u\Delta^{(m)} x$ for all $x \in \omega$ . In this study, the operator $\Delta_u^{(m)} : \omega \to \omega$ was defined as follows:

$$\Delta^{(m)} x_k = \sum_{i=0}^{m} (-1)^i \binom{m}{i} x_{k-i}. \tag{10}$$

Let $\alpha$ be a proper fraction and $u \in \mathcal{U}$. We define the sequence spaces $c_0(\Gamma, \Delta^{(\alpha)}, u)$ and $c(\Gamma, \Delta^{(\alpha)}, u)$ as follows:

$$c_0(\Gamma, \Delta^{(\alpha)}, u) = \{x \in \omega : (\sum_{j=0}^{k} u_j \Delta^{(\alpha)} x_j) \in c_0\} \tag{11}$$

and

$$c(\Gamma, \Delta^{(\alpha)}, u) = \{x \in \omega : (\sum_{j=0}^{k} u_j \Delta^{(\alpha)} x_j) \in c\}. \tag{12}$$

Now, we define the triangle matrix $\Delta_u^{(\alpha)}(\Gamma) = (\tau_{nk})$,

$$\tau_{nk} = \begin{cases} \sum_{i=0}^{n-k} (-1)^i \dfrac{\Gamma(\alpha+1)}{i!\Gamma(\alpha+1-i)} u_{i+k}, & (0 \leq k \leq n) \\ 0, & k > n \end{cases} \tag{13}$$

for all $k, n \in \mathbb{N}$. Further, for any sequence $x = (x_k)$ we define the sequence $y = (y_k)$ which will be used, as the $\Delta_u^{(\alpha)}(\Gamma)-$transform of $x$, that is

$$\begin{aligned} y_k &= \sum_{j=0}^{k} u_j \Delta^{(\alpha)} x_j = \sum_{j=0}^{k} u_j (x_j - \alpha x_{j-1} + \frac{\alpha(\alpha-1)}{2!} x_{j-3} + ...) \\ &= \sum_{j=0}^{k} \Big( \sum_{i=0}^{k-j} (-1)^i \frac{\Gamma(\alpha+1)}{i!\Gamma(\alpha+1-i)} u_{i+j} \Big) x_j \end{aligned} \tag{14}$$

for all $k \in \mathbb{N}$.

It is natural that the spaces $c_0(\Gamma, \Delta^{(\alpha)}, u)$ and $c(\Gamma, \Delta^{(\alpha)}, u)$ may also be defined with the notation of (2) that

$$c_0(\Gamma, \Delta^{(\alpha)}, u) = (c_0)_{\Delta_u^{(\alpha)}(\Gamma)} \quad \text{and} \quad c(\Gamma, \Delta^{(\alpha)}, u) = c_{\Delta_u^{(\alpha)}(\Gamma)}. \tag{15}$$

Before the main result let us give some lemmas, which we use frequently throughout this study, with respect to $\Delta^{(\alpha)}$ operator.

**Lemma 1.** *[20, Theorem 2.2]*

$$\Delta^{(\alpha)} o \Delta^{(\beta)} = \Delta^{(\beta)} o \Delta^{(\alpha)} = \Delta^{(\alpha+\beta)}.$$

**Lemma 2.** *[20, Theorem 2.3]*

$$\Delta^{(\alpha)} o \Delta^{(-\alpha)} = \Delta^{(-\alpha)} o \Delta^{(\alpha)} = Id$$

*where $Id$ the identity operator on $\omega$.*

**Theorem 1.** *The sequence spaces $c_0(\Gamma, \Delta^{(\alpha)}, u)$ and $c(\Gamma, \Delta^{(\alpha)}, u)$ are $BK-$spaces with the norm*

$$\|x\|_{c_0(\Gamma,\Delta^{(\alpha)},u)} = \|x\|_{c(\Gamma,\Delta^{(\alpha)},u)} = \sup_k \Big| \sum_{j=0}^{k} u_j \Delta^{(\alpha)} x_j \Big|. \tag{16}$$

PROOF. Since (15) holds and $c_0, c$ are $BK-$ spaces with respect to their natural norms (see [21, pp. 16-17]) and the matrix $\Delta_u^{(\alpha)}(\Gamma) = (\tau_{nk})$ is a triangle, Theorem 4.3.12 Wilansky [24, pp. 63] gives the fact that $c_0(\Gamma, \Delta^{(\alpha)}, u)$ and $c(\Gamma, \Delta^{(\alpha)}, u)$ are $BK-$ spaces with the given norms. This completes the proof .

Now, we may give the following theorem concerning the isomorphism between the spaces $c_0(\Gamma, \Delta^{(\alpha)}, u)$, $c(\Gamma, \Delta^{(\alpha)}, u)$ and $c_0, c$, respectively:

**Theorem 2.** *The sequence spaces $c_0(\Gamma, \Delta^{(\alpha)}, u)$ and $c(\Gamma, \Delta^{(\alpha)}, u)$ are linearly isomorphic to the spaces $c_0$ and $c$, respectively, i.e, $c_0(\Gamma, \Delta^{(\alpha)}, u) \cong c_0$ and $c(\Gamma, \Delta^{(\alpha)}, u) \cong c$.*

PROOF. We prove the theorem for the space $c(\Gamma, \Delta^{(\alpha)}, u)$. To prove this, we should show the existence of a linear bijection between the spaces $c(\Gamma, \Delta^{(\alpha)}, u)$ and $c$. Consider the transformation $T$ defined, with the notation of (14), from $c(\Gamma, \Delta^{(\alpha)}, u)$ to $c$ by $x \mapsto y = Tx = \Delta_u^{(\alpha)}(\Gamma)x$. The linearity of $T$ is clear. Further, it is trivial that $x = \theta$ whenever $Tx = \theta$ and hence $T$ is injective. Let be $y = (y_k) \in c$ and we define a sequence $x = (x_k) \in c(\Gamma, \Delta^{(\alpha)}, u)$ by

$$x_k = \sum_{i=0}^{\infty} (-1)^i \frac{\Gamma(1-\alpha)}{i!\Gamma(1-\alpha-i)} \frac{y_{k-i} - y_{k-i-1}}{u_{k-i}}. \tag{17}$$

Then by Lemma 2, we deduce that

$$\begin{aligned} \sum_{j=0}^{k} u_j \Delta^{(\alpha)} x_j &= \sum_{j=0}^{k} u_j \Delta^{(\alpha)} \Big( \sum_{i=0}^{\infty} (-1)^i \frac{\Gamma(1-\alpha)}{i!\Gamma(1-\alpha-i)} \frac{y_{j-i} - y_{j-i-1}}{u_{j-i}} \Big) \\ &= \sum_{j=0}^{k} u_j \Delta^{(\alpha)} \Big( \Delta^{(-\alpha)} \Big( \frac{y_j - y_{j-1}}{u_j} \Big) \Big) \\ &= \sum_{j=0}^{k} (y_j - y_{j-1}) \;=\; y_k \end{aligned} \tag{18}$$

Hence, $x \in c(\Gamma, \Delta^{(\alpha)}, u)$ so $T$ is surjective. Furthermore one can easily show that $T$ is norm preserving. This complete the proof.

## 3. The $\beta$-Dual of The Spaces $c_0(\Gamma, \Delta^{(\alpha)}, u)$ and $c(\Gamma, \Delta^{(\alpha)}, u)$

In this section, we determine the $\beta$-dual of the spaces $c_0(\Gamma, \Delta^{(\alpha)}, u)$ and $c(\Gamma, \Delta^{(\alpha)}, u)$. For the sequence spaces $X$ and $Y$, define the set $S(X, Y)$ by

$$S(X, Y) = \{z = (z_k) \in \omega : xz = (x_k z_k) \in Y \text{ for all } x \in X\}. \tag{19}$$

With the notation of (19), $\beta-$ dual of a sequence space $X$ is defined by

$$X^\beta = S(X, cs).$$

**Lemma 3.** *$A \in (c_0 : c)$ if and only if*

$$\lim_{n\to\infty} a_{nk} = \alpha_k \text{ for each fixed } k \in \mathbb{N}, \tag{20}$$

$$\sup_{n\in\mathbb{N}} \sum_k |a_{nk}| < \infty. \tag{21}$$

**Lemma 4.** *$A \in (c : c)$ if and only if (20) and (21) hold, and*

$$\lim_{n\to\infty} \sum_k a_{nk} \quad \text{exists.} \tag{22}$$

**Lemma 5.** *$A = (a_{nk}) \in (\ell_\infty : \ell_\infty)$ if and only if*

$$\sup_n \sum_k |a_{nk}| < \infty. \tag{23}$$

**Theorem 3.** *Define the sets $\Gamma_1$, $\Gamma_2$ and a matrix $T = (t_{nk})$ by*

$$t_{nk} = \begin{cases} t_k - t_{k+1}, & (k < n) \\ t_n, & (k = n) \\ 0, & (k > n) \end{cases}$$

*for all* $k,n\in\mathbb{N}$ *where* $t_k=a_k\sum_{i=0}^{k}(-1)^i\frac{\Gamma(1-\alpha)}{i!\Gamma(1-\alpha-i)}\frac{1}{u_{k-i}}$

$$\begin{aligned}\Gamma_1 &= \Big\{a=(a_n)\in w:\lim_{n\to\infty}t_{nk}=\alpha_k \text{ exists for each } k\in\mathbb{N}\Big\}\\ \Gamma_2 &= \Big\{a=(a_n)\in w:\sup_{n\in\mathbb{N}}\sum_k|t_{nk}|<\infty\Big\}\\ \Gamma_3 &= \Big\{a=(a_n)\in w:\sup_{n\in\mathbb{N}}\lim_{n\to\infty}\sum_k t_{nk} \quad \text{exists}\Big\}.\end{aligned}$$

*Then,* $\{c_0(\Gamma,\Delta^{(\alpha)},u)\}^\beta=\Gamma_1\cap\Gamma_2$ *and* $\{c(\Gamma,\Delta^{(\alpha)},u)\}^\beta=\Gamma_1\cap\Gamma_2\cap\Gamma_3$.

Proof. We prove the theorem for the space $c_0(\Gamma,\Delta^{(\alpha)},u)$. Let $a=(a_n)\in w$ and $x=(x_k)\in c_0(\Gamma,\Delta^{(\alpha)},u)$. Then, we obtain the equality

$$\begin{aligned}\sum_{k=0}^{n}a_kx_k &= \sum_{k=0}^{n}\Big[a_k\sum_{i=0}^{\infty}(-1)^i\frac{\Gamma(1-\alpha)}{i!\Gamma(1-\alpha-i)}\frac{1}{u_{k-i}}\Big](y_{k-i}-y_{k-i-1})\\ &= \sum_{k=0}^{n}\Big[a_k\sum_{i=0}^{k}(-1)^i\frac{\Gamma(1-\alpha)}{i!\Gamma(1-\alpha-i)}\frac{1}{u_{k-i}}\Big](y_{k-i}-y_{k-i-1})\\ &= \sum_{k=0}^{n-1}\Big[a_k\sum_{i=0}^{k}(-1)^i\frac{\Gamma(1-\alpha)}{i!\Gamma(1-\alpha-i)}\frac{1}{u_{k-i}}-a_{k+1}\sum_{i=0}^{k+1}(-1)^i\frac{\Gamma(1-\alpha)}{i!\Gamma(1-\alpha-i)}\frac{1}{u_{k+1-i}}\Big]y_k\\ &\quad+\Big[a_n\sum_{i=0}^{n}(-1)^i\frac{\Gamma(1-\alpha)}{i!\Gamma(1-\alpha-i)}\frac{1}{u_{n-i}}\Big]y_n\\ &= T_ny \qquad (24)\end{aligned}$$

Then we deduce by (24) that $ax=(a_kx_k)\in cs$ whenever $x=(x_k)\in c_0(\Gamma,\Delta^{(\alpha)},u)$ if and only if $Ty\in c$ whenever $y=(y_k)\in c_0$. This means that $a=(a_k)\in\{c_0(\Gamma,\Delta^{(\alpha)},u)\}^\beta$ if and only if $T\in(c_0:c)$. Therefore, by using Lemma 3, we obtain :

$$\lim_{n\to\infty}t_{nk}=\alpha_k \qquad \text{exists for each } k\in\mathbb{N}, \qquad (25)$$

$$\sup_{n\in\mathbb{N}}\sum_k|t_{nk}|<\infty. \qquad (26)$$

Hence, we conclude that $\{c_0(\Gamma,\Delta^{(\alpha)},u)\}^\beta=\Gamma_1\cap\Gamma_2$.

## 4. Some matrix transformations related to the sequence spaces $c_0(\Gamma, \Delta^{(\alpha)}, u)$ and $c(\Gamma, \Delta^{(\alpha)}, u)$

In this final section, we state some results which characterize various matrix mappings on the spaces $c_0(\Gamma, \Delta^{(\alpha)}, u)$ and $c(\Gamma, \Delta^{(\alpha)}, u)$.

We shall write throughout for brevity that

$$\widetilde{a}_{nk} = z_{nk} - z_{n,k+1} \quad \text{and} \quad b_{nk} = \sum_{j=0}^{n} \Big( \sum_{i=0}^{n-j} (-1)^i \frac{\Gamma(\alpha+1)}{i!\Gamma(\alpha+1-i)} u_{i+j} \Big) a_{jk} \tag{27}$$

for all $k, n \in \mathbb{N}$ , where

$$z_{nk} = a_{nk} \sum_{i=0}^{k} (-1)^i \frac{\Gamma(1-\alpha)}{i!\Gamma(1-\alpha-i)} \frac{1}{u_{k-i}}.$$

Now, we may give the following theorem.

**Theorem 4.** *Let $\lambda$ be any given sequence space and $\mu \in \{c_0, c\}$. Then, $A = (a_{nk}) \in (\mu(\Gamma, \Delta^{(\alpha)}, u) : \lambda)$ if and only if $C \in (\mu : \lambda)$ and*

$$C^{(n)} \in (\mu : c) \tag{28}$$

*for every fixed $n \in \mathbb{N}$, where $c_{nk} = \widetilde{a}_{nk}$ and $C^{(n)} = (c^{(n)}_{mk})$ with*

$$c^{(n)}_{mk} = \begin{cases} z_{nk} - z_{n,k+1}, & (k < m) \\ z_{nm}, & (k = m) \\ 0, & (k > m) \end{cases}$$

*for all $k, m \in \mathbb{N}$.*

Proof. Let $\lambda$ be any given sequence space. Suppose that (27) holds between the entries of $A = (a_{nk})$ and $C = (c_{nk})$, and take into account that the spaces $\mu(\Gamma, \Delta^{(\alpha)}, u)$ and $\mu$ are linearly isomorphic.

Let $A = (a_{nk}) \in (\mu(\Gamma, \Delta^{(\alpha)}, u) : \lambda)$ and take any $y = (y_k) \in \mu$. Then, $C\Delta^{(\alpha)}_u(\Gamma)$ exists and $\{a_{nk}\}_{k\in\mathbb{N}} \in \mu(\Gamma, \Delta^{(\alpha)}, u)^\beta$ which yields that (28) is necessary and $\{c_{nk}\}_{k\in\mathbb{N}} \in \mu^\beta$ for each $n \in \mathbb{N}$. Hence, $Cy$ exists for each $y \in \mu$ and thus by letting $m \to \infty$ in the equality

$$\sum_{k=0}^{m} a_{nk} x_k = \sum_{k=0}^{m-1} (z_{nk} - z_{n,k+1}) y_k + z_{nm} y_m; \quad (m, n \in \mathbb{N}) \tag{29}$$

we have that $Cy = Ax$ and so we have that $C \in (\mu : \lambda)$.

Conversely, suppose that $C \in (\mu : \lambda)$ and (28) hold, and take any $x = (x_k) \in \mu(\Gamma, \Delta^{(\alpha)}, u)$. Then, we have $\{c_{nk}\}_{k\in\mathbb{N}} \in \mu^\beta$ which gives together with (28) that $\{a_{nk}\}_{k\in\mathbb{N}} \in \mu(\Gamma, \Delta^{(\alpha)}, u)^\beta$ for each $n \in \mathbb{N}$. So, $Ax$ exists. Therefore, we obtain from the equality

$$\sum_{k=0}^{m} c_{nk} y_k = \sum_{k=0}^{m} \Big[ \sum_{j=k}^{m} \Big( \sum_{i=0}^{j-k} (-1)^i \frac{\Gamma(\alpha+1)}{i!\Gamma(\alpha+1-i)} u_{i+k} \Big) c_{nj} \Big] x_k \quad \text{for all } n \in \mathbb{N}, \tag{30}$$

as $m \to \infty$ that $Ax = Cy$ and this shows that $A \in (\mu(\Gamma, \Delta^{(\alpha)}, u) : \lambda)$. This completes the proof.

**Theorem 5.** *Suppose that the entries of the infinite matrices $A = (a_{nk})$ and $B = (b_{nk})$ are connected with the relation (27) and $\lambda$ be given sequence space and $\mu \in \{c_0, c\}$. Then $A = (a_{nk}) \in (\lambda : \mu(\Gamma, \Delta^{(\alpha)}, u))$ if and only if $B = (b_{nk}) \in (\lambda : \mu)$ .*

Proof. Let $z = (z_k) \in \lambda$ and consider the following equality with (27)

$$\sum_{k=0}^{m} b_{nk} z_k = \sum_{j=0}^{n} \Big[ \sum_{k=0}^{m} \Big( \sum_{i=0}^{n-j} (-1)^i \frac{\Gamma(\alpha+1)}{i!\Gamma(\alpha+1-i)} u_{i+j} \Big) a_{jk} \Big] z_k \quad (m, n \in \mathbb{N}),$$

which yields as $m \to \infty$ that $(Bz)_n = [\Delta_u^{(\alpha)}(\Gamma)(Az)]_n$. Hence, we obtain that $Az \in \mu(\Gamma, \Delta^{(\alpha)}, u)$ whenever $z \in \lambda$ if and only if $Bz \in \mu$ whenever $z \in \lambda$.

We will have several consequences by using Theorem 4 and Theorem 5. But we must give firstly some relations which are important for consequences:

$$\sup_{n\in\mathbb{N}} \sum_k |a_{nk}| < \infty \tag{31}$$

$$\lim_{n\to\infty} a_{nk} = \alpha_k \quad \text{exists for each fixed } k \in \mathbb{N} \tag{32}$$

$$\lim_{k\to\infty} a_{nk} = 0 \quad \text{for each fixed } n \in \mathbb{N} \tag{33}$$

$$\lim_{n\to\infty} \sum_k a_{nk} \quad \text{exists} \tag{34}$$

$$\lim_{n\to\infty} \sum_k a_{nk} = 0 \tag{35}$$

$$\sup_{K\in\mathcal{F}}\sum_{n}\Big|\sum_{k\in K}a_{nk}\Big|<\infty \tag{36}$$

$$\lim_{n\to\infty}\sum_{k}|a_{nk}|=0 \tag{37}$$

$$\sup_{n,k}|a_{nk}|<\infty \tag{38}$$

$$\lim_{m\to\infty}\sum_{k}|a_{nk}|=\sum_{k}|\alpha_k| \tag{39}$$

Now, we can give the corollaries:

**Corollary 1.** *The following statements hold:*

*(i)* $A=(a_{nk})\in(c_0(\Gamma,\Delta^{(\alpha)},u):\ell_\infty)=(c(\Gamma,\Delta^{(\alpha)},u):\ell_\infty)$ *if and only if (31) holds with* $\tilde{a}_{nk}$ *instead of* $a_{nk}$ *and (28) also holds.*

*(ii)* $A=(a_{nk})\in(c_0(\Gamma,\Delta^{(\alpha)},u):c)$ *if and only if (31)and (32) hold with* $\tilde{a}_{nk}$ *instead of* $a_{nk}$ *and (28) also holds.*

*(iii)* $A=(a_{nk})\in(c_0(\Gamma,\Delta^{(\alpha)},u):c_0)$ *if and only if (31) and (33) hold with* $\tilde{a}_{nk}$ *instead of* $a_{nk}$ *and (28) also holds.*

*(iv)* $A=(a_{nk})\in(c(\Gamma,\Delta^{(\alpha)},u):c)$ *if and only if (31), (32) and (34) hold with* $\tilde{a}_{nk}$ *instead of* $a_{nk}$ *and (28) also holds.*

*(v)* $A=(a_{nk})\in(c(\Gamma,\Delta^{(\alpha)},u):c_0)$ *if and only if (31), (33) and (35) hold with* $\tilde{a}_{nk}$ *instead of* $a_{nk}$ *and (28) also holds.*

*(vi)* $A=(a_{nk})\in(c_0(\Gamma,\Delta^{(\alpha)},u):\ell)=(c(\Gamma,\Delta^{(\alpha)},u):\ell)$ *if and only if (36) holds with* $\tilde{a}_{nk}$ *instead of* $a_{nk}$ *and (28) also holds.*

**Corollary 2.** *The following statements hold:*

*(i)*$A=(a_{nk})\in(\ell_\infty:c_0(\Gamma,\Delta^{(\alpha)},u))$ *if and only if (37) hold with* $b_{nk}$ *instead of* $a_{nk}$*.*

*(ii)*$A=(a_{nk})\in(c:c_0(\Gamma,\Delta^{(\alpha)},u))$ *if and only if (31), (33) and (35) hold with* $b_{nk}$ *instead of* $a_{nk}$*.*

*(iii)* $A=(a_{nk})\in(c_0:c_0(\Gamma,\Delta^{(\alpha)},u))$ *if and only if (31) and (33) hold with* $b_{nk}$ *instead of* $a_{nk}$*.*

*(iv)*$A=(a_{nk})\in(c:c_0(\Gamma,\Delta^{(\alpha)},u))$ *if and only if (33) and (38) hold with* $b_{nk}$ *instead of* $a_{nk}$*.*

*(v)* $A=(a_{nk})\in(\ell_\infty:c(\Gamma,\Delta^{(\alpha)},u))$ *if and only if (32) and (39) hold with* $b_{nk}$ *instead of* $a_{nk}$*.*

*(vi)* $A=(a_{nk})\in(c:c(\Gamma,\Delta^{(\alpha)},u))$ *if and only if (31), (32) and (34) hold with* $b_{nk}$ *instead of* $a_{nk}$*.*

*(vii)* $A = (a_{nk}) \in (c_0 : c(\Gamma, \Delta^{(\alpha)}, u))$ *if and only if (31) and (32) hold with* $b_{nk}$ *instead of* $a_{nk}$.

*(viii)* $A = (a_{nk}) \in (\ell : c(\Gamma, \Delta^{(\alpha)}, u))$ *if and only if (32) and (38) hold with* $b_{nk}$ *instead of* $a_{nk}$.

## References


[1] G.G. Lorentz, A contribution to the theory of divergent sequences, Acta Math. **80**(1948), 167-190.

[2] H. Kızmaz,On certain sequence spaces, Canad. Math. bull., 24(1981) 169-176

[3] M. Et, On some difference sequence spaces, Turkish J. Math. 17 (1993) 18-24.

[4] M. Et, R. Çolak, On some generalized difference sequence spaces, Soochow J. Math. 21 (1995) 377-386.

[5] R. Çolak, M. Et, E. Malkowsky, Some topics of sequence spaces, in: Lecture Notes in Mathematics, Fırat Univ. Press, Turkey, 2004, pp.1-63. Fırat Univ. Elazığ, 2004, ISBN:975-394-038-6.

[6] C. S. Wang, On Nörlund sequence spaces, Tamkang J. Math. **9**(1978), 269-274.

[7] P. N. Ng, P.Y. Lee, Cesàro sequence spaces of non-absolute type, Comment. Math. Prace Mat. **20**(2)(1978), 429-433.

[8] A. M. Jarrah, E. Malkowsky, BK-spaces, bases and linear operators, Ren. Circ. Mat. Palermo II **52**(1990), 177-191.

[9] E. Malkowsky, M. Mursaleen, S. Suantai, The dual spaces of sets of difference sequences of order $m$ and matrix transformations, Acta Math. Sin. Eng. Ser. 23(3)(2007) 521-532.

[10] B. Altay, On the space of $p-$ summable difference sequences of order $m$, $(1 \leq p < \infty)$, Studia Sci. Math. Hungar. 43(4)(2006) 387-402.

[11] B. Altay, F. Başar, Generalization of the space $\ell(p)$ derived by weighted mean , J. Math. Anal. Appl 330 (2007) 174-185.

[12] B. Altay, F. Başar, Certain topological properties and duals of the matrix domain of a triangle matrix in a sequence space, J. Math. Anal. Appl. 336(1)(2007) 632-645.

[13] M. Başarır, E. E. Kara, On the $m^{th}$ order difference sequence space of generalized weighted mean and compact operators, Acta Mathematica Scientia, 33(B3)(2013), 1-18.

[14] M. Başarır, E. E. Kara, On some difference sequence space of weighted means and compact operators, Annals of Functional Analysis, 2(2)(2011), 116-131.

[15] S. Demiriz, C. Çakan, Some topological and geometrical properties of a new difference sequence space, Abstract and Applied Analysis, In Press.

[16] Duyar O., Demiriz, S., On some new generalized difference sequence spaces and their topological properties, Journal of New Result in Science, 6(2014) 1-14.

[17] C. Aydın, F.Başar, Some new difference sequence spaces, Applied Mathematics and Computation, **157**(2004), 677-693.

[18] P. Baliarsingh, Some new difference sequence spaces of fractional order and their dual spaces, Applied Mathematics and Computation, 219(2013) 9737-9742.

[19] S. Dutta, P. Baliarsingh, A note on paranormed difference sequence spaces of fractional order and their matrix transformations, Journal of the Egyptian Mathematical Society, (2013).

[20] P. Baliarsingh, S. Dutta, A unifying approach to the difference operators and their applications, Bol. Soc. Paran. Mat., (3s)**v. 33** 1 (2015): 49-57.

[21] F. Başar, Summability Theory and Its Appliactions, Bentham Science Publishers, ISBN:978-1-60805-252-3, 2011.

[22] M. Stieglitz, H. Tietz, Matrix transformationen von folgenräumen eine ergebnisübersicht, Math. Z. 154(1977) 1-16.

[23] I.J. Maddox, Elements of Functional Analysis, second ed., The Cambridge University Press, Cambridge, 1988.

[24] A. Wilansky, Summability through Functional Analysis, North-Holland Mathematics Studies , Amsterdam, **85** 1984.